

# Sliding Mode Control for Integrator Systems

**part 3:** Noncontinuous Control Theory

# Contents

Bibliography .....	2
1 Discontinuous System Theory .....	3
1.1 Ternary Differential Equations' Solutions .....	4
1.2 Conditions for Existence and Uniqueness of Classical, Caratheodory, Filippov Solutions .....	5

# Bibliography

CORTES J, 2008. Discontinuous Dynamical Systems[J/OL]. IEEE Control Systems Magazine, 28(3): 36-73. DOI:10.1109/MCS.2008.919306.

# 1 Discontinuous System Theory

see (CORTES, 2008)

# 1.1 Ternary Differential Equations' Solutions

Table 1: solutions to ternary differential equations

differetial equation	differetial inclusion	classical solution	caratheodory solution	Filippov solution
$\dot{x} = \begin{cases} 1 & \text{if } x < 0 \\ a & \text{if } x = 0 \\ -1 & \text{if } x > 0 \end{cases}$	$\dot{x} \in \mathcal{F}(x) = \begin{cases} 1 & \text{if } x < 0 \\ [-1, 1] & \text{if } x = 0 \\ -1 & \text{if } x > 0 \end{cases}$	<p>Only when <math>a = 0</math>, classical solution exists. The maximal classical solution is</p> <ol style="list-style-type: none"> <li>1. if <math>x(0) &gt; 0, x_1(t) = x(0) - t, t &lt; x(0)</math></li> <li>2. if <math>x(0) &lt; 0, x_2(t) = x(0) + t, t &lt; -x(0)</math></li> <li>3. if <math>x(0) = 0, x_3(t) = 0, t \in [0, \infty)</math></li> </ol>	<p>Only when <math>a = 0</math>, caratheodory solution exists. The maximal classical solution is</p> <ol style="list-style-type: none"> <li>1. if <math>x(0) &gt; 0, x_1(t) = \max(x(0) - t, 0), t \in [0, \infty)</math></li> <li>2. if <math>x(0) &lt; 0, x_2(t) = \min(x(0) + t, 0), t \in [0, \infty)</math></li> <li>3. if <math>x(0) = 0, x_3(t) = 0, t \in [0, \infty)</math></li> </ol> <p><b>Note:</b> These only absolutely continuous (not continuously differentiable)</p>	<p>Whatever the value of <math>a</math> is, the Filippov solution is</p> <ol style="list-style-type: none"> <li>1. if <math>x(0) &gt; 0, x_1(t) = \max(x(0) - t, 0), t \in [0, \infty)</math></li> <li>2. if <math>x(0) &lt; 0, x_2(t) = \min(x(0) + t, 0), t \in [0, \infty)</math></li> <li>3. if <math>x(0) = 0, x_3(t) = 0, t \in [0, \infty)</math></li> </ol>
$\dot{x} = \begin{cases} -1 & \text{if } x < 0 \\ a & \text{if } x = 0 \\ 1 & \text{if } x > 0 \end{cases}$	$\dot{x} \in \mathcal{F}(x) = \begin{cases} -1 & \text{if } x < 0 \\ [-1, 1] & \text{if } x = 0 \\ 1 & \text{if } x > 0 \end{cases}$	<p>From <math>x = x(0) \neq 0</math>, classical solution exists as</p> <ol style="list-style-type: none"> <li>1. <math>x_1(t) = x(0) + t</math> if <math>x(0) &gt; 0</math></li> <li>2. <math>x_2(t) = x(0) - t</math> if <math>x(0) &lt; 0</math></li> </ol> <p>From <math>x = x(0) = 0</math>, classical solution exists when <math>a = 1</math> or <math>a = -1</math></p> <ol style="list-style-type: none"> <li>1. when <math>a = 1, x_1(t) = t, t \in [0, \infty)</math></li> <li>2. when <math>a = -1, x_2(t) = -t, t \in [0, \infty)</math></li> </ol>	<p>From <math>x = x(0) \neq 0</math>, classical solution exists as</p> <ol style="list-style-type: none"> <li>1. <math>x_1(t) = x(0) + t</math> if <math>x(0) &gt; 0</math></li> <li>2. <math>x_2(t) = x(0) - t</math> if <math>x(0) &lt; 0</math>.</li> </ol> <p>From <math>x = x(0) = 0</math>, two caratheodory solutions exist for <b>all</b> <math>a \in \mathbb{R}</math></p> <ol style="list-style-type: none"> <li>1. <math>x_1(t) = t, t \in [0, \infty)</math></li> <li>2. <math>x_2(t) = -t, t \in [0, \infty)</math></li> </ol> <p>These two solutions only violate the vector field in <math>t = 0</math></p>	<p>Filippov solution exists for all <math>a \in \mathbb{R}</math> and <math>x(0) \in \mathbb{R}</math>.</p> <ol style="list-style-type: none"> <li>1. if <math>x(0) \geq 0, x_1(t) = x(0) + t, t \in [0, \infty)</math></li> <li>2. if <math>x(0) \leq 0, x_2(t) = x(0) - t, t \in [0, \infty)</math></li> </ol> <p><b>Note:</b> When <math>x(0) = 0</math>, exists two Filippov solutions.</p>
$\dot{x} = \begin{cases} 1 & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$	$\dot{x} \in \{1\}$	$x = 0, t \in [0, \infty)$	<p>two caratheodory solutions:</p> <ol style="list-style-type: none"> <li>1. <math>x(t) = 0, t \in [0, \infty)</math></li> <li>2. <math>x(t) = t, t \in [0, \infty)</math></li> </ol>	<p>one unique solution:</p> <ol style="list-style-type: none"> <li>1. <math>x(t) = t, t \in [0, \infty)</math></li> </ol>

## 1.2 Conditions for Existence and Uniqueness of Classical, Caratheodory, Filippov Solutions

Table 2: conditions of solutions to  $\dot{x} = X(x(t))$

	solution	existence	uniqueness
classical	continuously differentiable	$X : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is continuous	essentially one-sided Lipschitz on $B(x, \varepsilon)$ , <sup>1</sup>
Filippov	absolutely continuous	$X : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is measurable and locally essentially bounded	essentially one-sided Lipschitz on $B(x, \varepsilon)$

---

<sup>1</sup>Every vector field that is locally Lipschitz at  $x$  satisfies the one-sided Lipschitz condition on a neighborhood of  $x$ , but the converse is not true.

# THANKS